

2021

## MATHEMATICS — HONOURS

Fifth Paper

(Module - IX)

Full Marks : 50

*The figures in the margin indicate full marks.**Candidates are required to give their answers in their own words as far as practicable.* $\mathbb{R}$ ,  $\mathbb{N}$  denote the set of real numbers and the set of natural numbers respectively.Answer **question no. 1** and **any four** questions from the rest.1. (a) Answer **any two** questions :(i) Prove or disprove :  $T = \left\{1 - \frac{1}{n^2} : n \in \mathbb{N}\right\}$  is compact. 2(ii) Correct or justify : If a real valued function  $f$  is bounded in some closed interval  $[a, b]$  in  $\mathbb{R}$  then  $f$  is a function of bounded variation in  $[a, b]$ . 2(iii) Correct or justify : The power series  $x + \frac{x^2}{2^2} + \frac{x^3}{3^3} + \frac{x^4}{4^4} \dots$  is everywhere convergent. 2(iv) Discuss the continuity of the limit function of the sequence of functions  $\{f_n\}_n$  defined by

$$f_n(x) = \frac{x^{2n}}{1 + x^{2n}} \text{ on } [0, 1]. \quad 2$$

(b) Answer **any two** questions :(i) Examine whether  $\lim_{x \rightarrow 0} \frac{\int_0^{x^2} e^{\sqrt{1+t}} dt}{x^2} = e$ . 3(ii) If  $f$  is differentiable on  $[0, 1]$ , then  $\int_0^1 f'(x) dx = f(1) - f(0)$ . 3(iii) Cite with justification an example of a function  $f$  such that  $\frac{1}{f}$  is Riemann integrable but  $f$  is not so over its domain. 3**Please Turn Over**

(iv) Let  $H = (0, 1) \subseteq \mathbb{R}$  and  $\mathcal{G} = \{I_x : x \in H\}$  where  $I_x = \left(\frac{x}{2}, \frac{x+1}{2}\right)$ . Verify whether  $\mathcal{G}$  is an open cover of  $H$ . 3

2. (a) If  $S$  is a bounded and closed set of real numbers, then prove that every infinite open cover of  $S$  has a finite subcover. 5

(b) Let  $T = \left\{x \in \mathbb{R} : \cos \frac{1}{x} = 0\right\} \cup \{0\}$ . Is  $\mathbb{R} \setminus T$  compact? Justify your answer. 2

(c) Examine whether the following function is of bounded variation :

$f : [0, 1] \rightarrow \mathbb{R}$  defined by  $f(x) = \begin{cases} x \sin \frac{\pi}{x}, & x \in (0, 1] \\ 0, & x = 0 \end{cases}$ . 3

3. (a) If two functions  $f$  and  $g$  are Riemann integrable on  $[a, b]$ , use Lebesgue's theorem to prove that  $|f| - fg$  is Riemann integrable on  $[a, b]$ . 4

(b) Let a function  $f : [0, 3] \rightarrow \mathbb{R}$  be defined by  $f(x) = \begin{cases} x & \text{for } 0 \leq x \leq 1 \\ 1 & \text{for } 1 < x \leq 2 \\ x-1 & \text{for } 2 < x \leq 3 \end{cases}$  and

let  $F(x) = \int_0^x f(t) dt$  for  $0 \leq x \leq 3$ . Verify whether  $F$  is derivable on  $[0, 3]$ . 3

(c) Let  $f$  and  $g$  be continuous functions on a closed interval  $[a, b]$  and  $\int_a^b f(x) dx = \int_a^b g(x) dx$ .

Show that there exist a point  $c \in [a, b]$  for which  $f(c) = g(c)$ . 3

4. (a) State and prove Darboux's Theorem on upper Riemann integral. 1+4

(b) Give example of a Riemann integrable function that has no primitive. 2

(c) Show that  $\left| \int_0^{\pi/2} \sin x \cos(x^2) dx \right| \leq \frac{1}{2}$  3

5. (a) Give examples (with justification) of Riemann integrable functions  $f, g$  on  $[0, 1]$  such

that  $\int_0^1 |f - g| = 0$ , but  $f \neq g$ . 3

- (b) Let  $f : [a, b] \rightarrow \mathbb{R}$  be Riemann integrable over  $[a, b]$  and  $g : [a, b] \rightarrow \mathbb{R}$  be a function such that 'g' differs from 'f' at finitely many points of  $[a, b]$ . Prove that  $g$  is also Riemann integrable over

$$[a, b] \quad \text{and} \quad \int_a^b f = \int_a^b g. \quad 2+2$$

- (c) Cite with justification an example of a function  $f$  such that  $|f|$  is Riemann integrable but  $f$  is not so over its domain. 3

6. (a) Examine the applicability of Weierstrass' form of Second Mean Value Theorem of Integral Calculus

$$\text{for } \int_0^{\pi} x^2 \sin x dx. \quad 3$$

- (b) State Dini's Theorem on sequence of real valued functions. If  $f_n(x) = x^n(1-x)$ , where  $\{f_n\}_n$  is a sequence of functions defined on  $[0, 1]$  then by using Dini's theorem, prove that  $f_n \rightarrow 0$  uniformly on  $[0, 1]$ . 1+3

- (c) A sequence of functions  $\{f_n\}_n$  is defined by  $f_n(x) = \sqrt{x^2 + \frac{1}{n^2}}$ , where  $x \in [-1, 1]$ . Show that  $\{f_n\}_n$  is uniformly convergent on  $[-1, 1]$ . 3

7. (a) State Dirichlet's test on uniform convergence for series of functions. Prove that the series  $\sum_{n=1}^{\infty} \frac{\sin nx}{n}$  is uniformly convergent on any closed interval  $[a, b]$  contained in the open interval  $(0, 2\pi)$ . 2+3

- (b) Correct or justify :

$$\text{If } \sum_{n=0}^{\infty} |a_n| \text{ is convergent then } \int_0^1 \left( \sum_{n=0}^{\infty} a_n x^n \right) dx = \sum_{n=0}^{\infty} \frac{a_n}{n+1} \quad 3$$

- (c) Prove or disprove : The function defined by  $f(x) = \sum_{n=1}^{\infty} \frac{\cos nx}{10^n}$ ,  $x \in \mathbb{R}$  is continuous everywhere. 2

8. (a) Find the radius of convergence of the power series  $x + \frac{(2!)^2 x^2}{4!} + \frac{(3!)^2 x^3}{6!} + \dots + \frac{(n!)^2 x^n}{(2n)!} + \dots$  3

- (b) Assuming the power series expansion for  $(1-x^2)^{-1/2}$  as

$$\frac{1}{\sqrt{1-x^2}} = 1 + \frac{1}{2}x^2 + \frac{1.3}{2.4}x^4 + \frac{1.3.5}{2.4.6}x^6 + \dots; |x| < 1.$$

Obtain the power series for  $\sin^{-1}x$  in  $(-1, 1)$ . 4

**Please Turn Over**

- (c) Correct or justify : If  $\sum_{n=0}^{\infty} a_n x^n$  converges at  $c \in \mathbb{R} \setminus \{0\}$ , then it converges absolutely for all  $x$  such that  $|x| < |c|$ .